arXiv:math/0010321v2 [math.QA] 5 Dec 2000

A PROOF OF THE TSYGAN FORMALITY CONJECTURE FOR CHAINS

BORIS SHOIKHET

Abstract. We extend the Kontsevich formality L_∞-morphism $\mathcal{U}\colon T^\bullet_{\mathrm{poly}}(\mathbb{R}^d) \to \mathcal{D}^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ to an L_∞-morphism of L_∞-modules over $T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$, $\hat{\mathcal{U}}\colon C_\bullet(A,A) \to \Omega^\bullet(\mathbb{R}^d)$, $A = C^\infty(\mathbb{R}^d)$. The construction of the map $\hat{\mathcal{U}}$ is given in Kontsevich-type integrals. The conjecture that such an L_∞-morphism exists is due to Boris Tsygan [Ts]. As an application, we obtain an explicit formula for isomorphism $A_*/[A_*,A_*] \xrightarrow{\sim} A/\{A,A\}$ (A_* is the Kontsevich deformation quantization of the algebra A by a Poisson bivector field, and $\{,\}$ is the Poisson bracket). We also formulate a conjecture extending the Kontsevich theorem on cup-products to this context. The conjecture implies a generalization of the Duflo formula, and many other things.

1. L_∞-ALGEBRAS AND L_∞-MODULES

Here we recall basic definitions from [Ts] and construct an L_∞-module over $T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ structure on the chain Hochschild complex $C_\bullet(A,A)$, $A = C^\infty(\mathbb{R}^d)$.

1.1. For a manifold M, $T^\bullet_{\mathrm{poly}}(M)$ denotes the graded Lie algebra of smooth polyvector fields on M, $\Omega^\bullet(M)$ denotes the graded space of smooth differential forms on M. The space $T^\bullet_{\mathrm{poly}}(M)$ is graded as a Lie algebra, i.e. $T^i_{\mathrm{poly}}(M) = \{(i+1)\text{-polyvector fields}\}$, $i \geq -1$. The bracket is the Schouten–Nijenhuis bracket, generalizing the usual Lie bracket of vector fields. It is defined as follows:

$$(1)\quad [\xi_0\wedge\dots\wedge\xi_k, \eta_0\wedge\dots\wedge\eta_l] = \\ = \sum_{i=0}^k\sum_{j=0}^l (-1)^{i+j+k}[\xi_i,\eta_j]\wedge\xi_0\wedge\dots\wedge\hat{\xi}_i\wedge\dots\wedge\xi_k\wedge\eta_0\wedge\dots\wedge\hat{\eta}_j\wedge\dots\wedge\eta_l$$

for $k,l \geq 0$ ($\{\xi_i\}$ and $\{\eta_j\}$ are vector fields, the bracket (1) does not depend on their choice and depends only on polyvector fields $\xi = \xi_0\wedge\dots\wedge\xi_k$ and $\eta = \eta_0\wedge\dots\wedge\eta_l$) and for $k \geq 0$

$$(2)\qquad [\xi_0\wedge\dots\wedge\xi_k, h] = \sum_{i=0}^k (-1)^i \xi_i(h)\xi_0\wedge\dots\wedge\hat{\xi}_i\wedge\dots\wedge\xi_k$$

(here $h \in C^\infty(M)$ is a function).

It is clear that $[T^i_{\mathrm{poly}}, T^j_{\mathrm{poly}}] \subset T^{i+j}_{\mathrm{poly}}$. We will consider $T^\bullet_{\mathrm{poly}}$ as a differential graded (dg) Lie algebra equipped with zero differential.

The space $\Omega^\bullet(M)$ is $\mathbb{Z}_{\leq 0}$-graded:

$$\Omega^i(M) = \{i\text{-differential forms}\},\quad \deg\Omega^i = -i.$$

Date: 2000.

For a k-polyvector field γ we denote by i_γ the natural contraction

$$i_\gamma\colon \Omega^\bullet(M) \to \Omega^{\bullet-k}(M).$$

The Lie derivative L_γ is defined by the Cartan formula

$$L_\gamma = d_{\mathrm{DR}} \circ i_\gamma + i_\gamma \circ d_{\mathrm{DR}} = [d_{\mathrm{DR}}, i_\gamma] \tag{3}$$

(we denote $[a,b]_+$ by $[a,b]$ for odd symbols a, b).

Lemma. $[L_{\gamma_1}, L_{\gamma_2}] = L_{[\gamma_1,\gamma_2]}$ *where* $[\gamma_1,\gamma_2]$ *is the Schouten–Nijenhuis bracket* (1), (2) *of polyvector fields.*

In this way we endow the graded space $\Omega^\bullet(M)$ with a structure of a module over the graded Lie algebra $T^\bullet_{\mathrm{poly}}(M)$. Note that for $\gamma \in T^k_{\mathrm{poly}}(M)$ one has $L_\gamma \Omega^\bullet(M) \subset \Omega^{\bullet-k}(M)$ and $\deg \Omega^{\bullet-k} = \deg \Omega^\bullet + k$ because we consider $\Omega^\bullet$ to be $\mathbb{Z}_{\le 0}$ graded.

1.2. We denote by $C^\bullet(A,A)$ the cohomological Hochschild complex of an associative algebra A. By definition, $C^k(A,A) = \mathrm{Hom}_{\mathbb{C}}(A^{\otimes k}, A)$, $k \ge 0$. The differential $d_{\mathrm{Hoch}}\colon C^k(A,A) \to C^{k+1}(A,A)$ is defined as follows:

$$\begin{aligned}(d\Psi)(a_1 \otimes \dots \otimes a_{k+1}) &= a_1 \cdot \Psi(a_2 \otimes \dots \otimes a_{k+1}) - \\ &- \Psi(a_1 \cdot a_2 \otimes a_3 \otimes \dots \otimes a_{k+1}) + \dots \pm \Psi(a_1 \otimes \dots \otimes a_k) \cdot a_{k+1}.\end{aligned} \tag{4}$$

By definition, $\mathcal{D}^\bullet_{\mathrm{poly}}(M) = C^\bullet_{\mathrm{poly}}(C^\infty(M), C^\infty(M))[1]$ is shifted by 1 subcomplex of $C^\bullet(A,A)$, $A = C^\infty(M)$, consisting of polydifferential operators. Thus, $\mathcal{D}^k_{\mathrm{poly}}(M) \subset \mathrm{Hom}_{\mathbb{C}}(A^{\otimes k+1}, A)$, $k \ge -1$.

The Gerstenhaber bracket on $C^\bullet(A,A)[1]$ is defined as follows: for $\phi_1 \in C^{k_1+1}(A,A)$, $\phi_2 \in C^{k_2+1}(A,A)$

$$[\phi_1, \phi_2] = \phi_1 \circ \phi_2 - (-1)^{k_1 k_2} \phi_2 \circ \phi_1 \tag{5}$$

where

$$\begin{aligned}(\phi_1 \circ \phi_2)(a_0 \otimes \dots \otimes a_{k_1+k_2}) &= \\ = \sum_{i=0}^{k_1} (-1)^{i k_2} \phi_1(a_0 \otimes \dots \otimes a_{i-1} \otimes \phi_2(a_i \otimes \dots \otimes a_{i+k_2}) &\otimes a_{i+k_2+1} \otimes \dots \otimes a_{k_1+k_2}).\end{aligned} \tag{6}$$

Lemma. *Formulas* (4), (5), (6) *define a dg Lie algebra structure on* $C^\bullet(A,A)[1]$.

Note that

$$d_{\mathrm{Hoch}}(\Psi) = [m, \Psi] \tag{7}$$

where $m\colon A^{\otimes 2} \to A$ is the product, the associativity is equivalent to $[m,m] = 0$, and $\mathrm{ad}\, m$ defines a differential.

1.2.1.

Lemma (Hochschild–Kostant–Rosenberg)**.**

(i) $H^i(\mathcal{D}^\bullet_{\mathrm{poly}}(M)) = T^i_{\mathrm{poly}}(M)$, *and the map* $\varphi_{\mathrm{HKR}}\colon T^\bullet_{\mathrm{poly}} \to \mathcal{D}^\bullet_{\mathrm{poly}}$,

$$\varphi_{\mathrm{HKR}}(\xi_1 \wedge \dots \wedge \xi_k)(f_1 \otimes \dots \otimes f_k) = \frac{1}{k!} \operatorname*{Alt}_{\xi_1,\dots,\xi_k} \prod_{i=1}^{k} \xi_i(f_i) \tag{8}$$

is a quasi-isomorphism of complexes (*in particular,* $d_{\mathrm{Hoch}}|_{\mathrm{Im}\,\varphi_{\mathrm{HKR}}} = 0$);

(ii) *the induced map on cohomology* $[\varphi_{\mathrm{HKR}}]\colon T^\bullet_{\mathrm{poly}}(M)\to H^\bullet(\mathcal{D}^\bullet_{\mathrm{poly}}(M))$ *is a Lie algebras isomorphism.*

Formality theorem of M. Kontsevich [K] states that $T^\bullet_{\mathrm{poly}}(M)$ and $\mathcal{D}^\bullet_{\mathrm{poly}}(M)$ are quasi-isomorphic as dg Lie algebras, in the sense of the derived categories, or L_∞-quasi-isomorphic.

1.3. We denote by $C_\bullet(A,A)$ the homological (chain) Hochschild complex of an associative algebra A. By definition, $C_k(A,A)=A\otimes A^{\otimes k}$, and the differential $b\colon C_k(A,A)\to C_{k-1}(A,A)$ is defined as follows:

$$b(a_0\otimes\dots\otimes a_k)= \\ =a_0a_1\otimes a_2\otimes\dots\otimes a_k-a_0\otimes a_1a_2\otimes\dots\otimes a_k+\dots\pm a_ka_0\otimes a_1\otimes\dots\otimes a_{k-1}. \tag{9}$$

We consider $C_\bullet(A,A)$ to be $\mathbb{Z}_{\le 0}$-graded, $\deg(A\otimes A^{\otimes k})=-k$, $k\ge 0$.

1.3.1.

Theorem.

(i) $H_i(C_\bullet(A,A))=\Omega^i(M)$, $A=C^\infty(M)$;

(ii) *the map*

$$\begin{gathered}\mu\colon C_\bullet(A,A)\to\Omega^\bullet(M),\\ \mu(a_0\otimes a_1\otimes\dots\otimes a_k)=\frac{1}{k!}a_0\,da_1\wedge\dots\wedge da_k\end{gathered} \tag{10}$$

is a quasi-isomorphism of the complexes ($\Omega^\bullet(M)$ *is equipped with zero differential*).

1.4. In Section 1.1 we have defined operators L_γ, $\gamma\in T^\bullet_{\mathrm{poly}}(M)$, acting on $\Omega^\bullet(M)$. We know from Theorems 1.2.1(i), 1.3.1(i) that $H^\bullet(\mathcal{D}^\bullet_{\mathrm{poly}}(M))=T^\bullet_{\mathrm{poly}}(M)$, $H_\bullet(C_\bullet(A,A))=\Omega^\bullet(M)$, $A=C^\infty(M)$. In this section we define the operators L_γ "on the level of complexes", i.e., we define operators L_Ψ, $\Psi\in C^\bullet(A,A)[1]$, acting on $C_\bullet(A,A)$. The operator L_Ψ, $\Psi\in\mathrm{Hom}(A^{\otimes k},A)$, is defined as follows:

(11)

$$L_\Psi(a_0\otimes\dots\otimes a_n)=\sum_{i=0}^{n-k}(-1)^{(k-1)(i+1)}a_0\otimes\dots\otimes a_i\otimes\Psi(a_{i+1}\otimes\dots\otimes a_{i+k})\otimes\dots\otimes a_n+ \\ +\sum_{j=n-k}^{n}(-1)^{n(j+1)}\Psi(a_{j+1}\otimes\dots\otimes a_0\otimes\dots)\otimes a_{k+j-n}\otimes\dots\otimes a_j.$$

Note that

$$L_m(a_0\otimes\dots\otimes a_n)=b(a_0\otimes\dots\otimes a_n) \tag{12}$$

(here $m\colon A^{\otimes 2}\to A$ is the multiplication, and b is the chain Hochschild differential, see (9)).

1.4.1.

Lemma.

$$[L_{\Psi_1}, L_{\Psi_2}] = L_{[\Psi_1,\Psi_2]}$$

where $[\Psi_1, \Psi_2]$ *is the Gerstenhaber bracket of the cochains* Ψ_1 *and* Ψ_2*.*

In particular, let $\Psi_1 = m\colon A^{\otimes 2} \to A$ be the multiplication; we know that $L_m = b$, and we have:

$$[b, L_\Psi] = L_{d_{\mathrm{Hoch}}\Psi} \tag{13}$$

(see (7)).

Lemma 1.4.1 allows us to equip the space $C_\bullet(A, A)$ with a structure of the module over the dg Lie algebra $C^\bullet(A, A)[1]$. We explain below how, using an L_∞-map $\mathcal{U}\colon T^\bullet_{\mathrm{poly}} \to \mathcal{D}^\bullet_{\mathrm{poly}}$, to equip the space $C_\bullet(A, A)$ with an L_∞-module structure over $T^\bullet_{\mathrm{poly}}$.

1.5. In this Subsection we recall the basic definitions of homotopical algebra. We will do it "on the level of formulas", leaving the language of formal Q-manifolds [K], because it is more convenient for our needs.

1.5.1. An L_∞-algebra is a $\mathbb{Z}$-graded vector space $\mathfrak{g}$ with a collection of maps:

$$\begin{aligned} &Q_1\colon \mathfrak{g} \to \mathfrak{g}[1] \\ &Q_2\colon \Lambda^2\mathfrak{g} \to \mathfrak{g} \\ &Q_3\colon \Lambda^3\mathfrak{g} \to \mathfrak{g}[-1] \\ &\dots\dots\dots\dots\dots\dots \end{aligned}$$

satisfying the relations

$$\sum_{\substack{i_1<\dots<i_p,\ j_1<\dots<j_q \\ p+q=k}} \pm Q_{q+1}(Q_p(x_{i_1} \wedge \dots \wedge x_{i_p}) \wedge x_{j_1} \wedge \dots \wedge x_{j_p}) = 0 \tag{14}$$

for each $k \geq 2$ and homogenous $\{x_s\}$.

The first relation, for $k = 2$, is

$$Q_1^2 = 0.$$

The second, $k = 3$, is that the product Q_2 is compatible with the differential Q_1, i.e.

$$Q_1(Q_2(x \wedge y)) = Q_2(Q_1 x \wedge y) \pm Q_2(x \wedge Q_1 y).$$

The third is that the skew-symmetric product Q_2 obeys the Jacobi identity modulo Q_3. The case when $Q_3 = Q_4 = \dots = 0$ is the case of dg Lie algebras.

1.5.2. An L_∞-morphism $\mathcal{U}\colon \mathfrak{g}_1 \to \mathfrak{g}_2$ between two L_∞-algebras is a collection of maps:

$$\begin{aligned} &\mathcal{U}_1\colon \mathfrak{g}_1 \to \mathfrak{g}_2 \\ &\mathcal{U}_2\colon \Lambda^2\mathfrak{g}_1 \to \mathfrak{g}_2[-1] \\ &\mathcal{U}_3\colon \Lambda^3\mathfrak{g}_1 \to \mathfrak{g}_2[-2] \\ &\dots\dots\dots\dots\dots\dots \end{aligned}$$

obeying the identities

$$\begin{aligned}\sum \pm \mathcal{U}_{q+1}(Q_p(x_{i_1}\wedge\cdots\wedge x_{i_p})\wedge x_{j_1}\wedge\cdots\wedge x_{j_q}) &= \\ = \sum \pm \frac{1}{k!} Q_k(\mathcal{U}_{n_1}(x_{i_{11}}\wedge\cdots\wedge x_{i_{1n_1}})\wedge\cdots\wedge\mathcal{U}_{n_k}(x_{i_{k1}}\wedge\cdots\wedge x_{i_{kn_k}}))& \\ (n_1+\cdots+n_k=p+q).&\end{aligned}\tag{15}$$

In the simplest case, when $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are dg Lie algebras, $Q_1=d$, $Q_2=[,]$, (15) is:

$$\begin{aligned}d\mathcal{U}_n(x_1\wedge\cdots\wedge x_n)+\sum_{i=1}^{n}\pm\mathcal{U}_n(x_1\wedge\cdots\wedge dx_i\wedge\cdots\wedge x_n) &= \\ =\frac{1}{2}\sum_{\substack{k,l\geq 1\\ k+l=n}}\frac{1}{k!l!}\sum_{\sigma\in\Sigma_n}\pm[\mathcal{U}_k(x_{\sigma_1}\wedge\cdots\wedge x_{\sigma_k}),\mathcal{U}_l(x_{\sigma_{k+1}}\wedge\cdots\wedge x_{\sigma_n})]+& \\ +\sum_{i<j}\pm\mathcal{U}_{n-1}([x_i,x_j]\wedge x_1\wedge\cdots\wedge\hat{x}_i\wedge\cdots\wedge\hat{x}_j\wedge\cdots\wedge x_n).&\end{aligned}\tag{16}$$

The simplest cases are:

- **$n=1$:** $d\mathcal{U}_1(x)=\mathcal{U}_1(dx)$, i.e. $\mathcal{U}_1$ is a map of complexes; an L_∞-morphism is called an L_∞-quasi-isomorphism if $\mathcal{U}_1$ is a quasi-isomorphism of complexes;
- **$n=2$:** $\mathcal{U}_1$ is a map of dg Lie algebras modulo $\mathcal{U}_2$; $\mathcal{U}_1$ is a map of graded Lie algebras on the level of cohomology.

The connection of the notion of an L_∞-morphism with the classical homological algebra is that if two dg Lie algebras $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are quasi-isomorphic in the sense of derived cathegeries, that is there exists a hat $\mathfrak{g}_1 \xrightarrow{\varphi_1} \mathfrak{g}_3 \xleftarrow{\varphi_2} \mathfrak{g}_2$ for some dg Lie algebra $\mathfrak{g}_3$, and both maps φ_1 and φ_2 are maps of dg Lie algebras and quasi-isomorphisms of the complexes, they are L_∞-quasi-isomorphic. In the L_∞-side we do not construct an extra dg Lie algebra $\mathfrak{g}_3$ but we construct an infinitely many higher "Taylor components" of the map $\mathcal{U}_1$.

1.5.3. An L_∞-module M over an L_∞-algebra $\mathfrak{g}$ is a graded vector space $\mathfrak{g}$ with a collection of maps

$$\varphi_k\colon \Lambda^k\mathfrak{g}\otimes M\to M[1-k],\quad k\geq 0$$

satisfying for each $k\geq 0$ the equation

$$\begin{aligned}\sum_{p+q=k}\sum_{\substack{i_1<\cdots<i_p\\ j_1<\cdots<i<j_q}}\pm\phi_p(x_{i_1}\wedge\cdots\wedge x_{i_p}\otimes\phi_q(x_{j_1}\wedge\cdots\wedge x_{j_q}\otimes m))+& \\ +\sum_{p+q=k}\sum_{\substack{i_1<\cdots<i_p\\ j_1<\cdots<j_q}}\pm\phi_{q+1}(Q_p(x_{i_1}\wedge\cdots\wedge x_{i_p})\wedge x_{j_1}\wedge\cdots\wedge x_{j_q}\otimes m)=0.&\end{aligned}\tag{17}$$

For $k=0$ (17) gives $\phi_0^2=0$, i.e. ϕ_0 is a differential on th graded space M.

For $k=1$ we obtain that the map $\phi_1\colon\mathfrak{g}\otimes M\to M$ is a map of the complexes:

$$\phi_0(\phi_1(x\otimes m))=\phi_1(dx\otimes m)\pm\phi_1(x\otimes\phi_0 m).$$

In the case when $\mathfrak{g}$ is a dg Lie algebra we obtain that the map ϕ_1 defines a $\mathfrak{g}$-module structure on M modulo ϕ_2, etc.

1.5.4. A morphism φ of two L_∞-modules M, N over an L_∞-algebra $\mathfrak{g}$ is a collection of maps

$$\varphi_k\colon \Lambda^k\mathfrak{g}\otimes M\to N[-k],\quad k\ge 0$$

satisfying

$$\begin{aligned}(18)\quad &\sum \pm\varphi_{q+1}(Q_p(x_{i_1}\wedge\dots\wedge x_{i_p})\wedge x_{j_1}\wedge\dots\wedge x_{j_q}\otimes m)+\\ &+\sum\pm\varphi_{p+1}(x_{i_1}\wedge\dots\wedge x_{i_p}\wedge\phi_q(x_{j_1}\wedge\dots\wedge x_{j_q}\otimes m))=\\ &=\sum\pm\phi_{p+1}(x_{i_1}\wedge\dots\wedge x_{i_p}\wedge\varphi_q(x_{j_1}\wedge\dots\wedge x_{j_q}\otimes m))\end{aligned}$$

for $p+q=k\ge 0$.

For $k=p+q=0$ we obtain that the map φ_0 is a map of the complexes, etc.

1.6. **An L_∞-module over $T^\bullet_{\text{poly}}(\mathbb{R}^d)$ structure on $C_\bullet(A,A)$.** M. Kontsevich constructed in [K] an L_∞-(quasi-iso)morphism $\mathcal{U}\colon T^\bullet_{\text{poly}}(\mathbb{R}^d)\to\mathcal{D}^\bullet_{\text{poly}}(\mathbb{R}^d)$. Let $\mathcal{U}_1,\mathcal{U}_2,\mathcal{U}_3,\dots$ be its Taylor components. Define a set of maps

$$\phi_k\colon\Lambda^k T^\bullet_{\text{poly}}(\mathbb{R}^d)\otimes C_\bullet(A,A)\to C_\bullet(A,A)[1-k]$$

by the formulas:

$$(19)\qquad \left.\begin{array}{l}\phi_0=b\ \text{(see (9))},\\ \phi_k(\gamma_1\wedge\dots\wedge\gamma_k\otimes\omega)=L_{\Psi_k}\omega\\ \ \text{with } \Psi_k=\mathcal{U}_k(\gamma_1\wedge\dots\wedge\gamma_k)\ \text{for } k\ge 1\end{array}\right\}$$

Lemma. *In this way we have defined an L_∞-module structure on $C_\bullet(A,A)$ over dg Lie algebra $T^\bullet_{\text{poly}}(\mathbb{R}^d)$.*

Proof. We have to prove (17) for $\{\phi_k\}$.

For $k=0$ it is just $b^2=0$,

for $k=1$ it is

$$(20)\qquad b\circ L_{\mathcal{U}_1(\gamma)}\omega=\pm L_{\mathcal{U}_1(\gamma)}\circ b\omega$$

But we have from (13):

$$[b,L_{\mathcal{U}_1(\gamma)}]\omega=L_{d_{\text{Hoch}}\mathcal{U}_1(\gamma)}\omega$$

and $d_{\text{Hoch}}\mathcal{U}_1(\gamma)=\mathcal{U}_1(d_{T_{\text{poly}}}\gamma)\equiv 0$ for any L_∞-morphism $\mathcal{U}\colon T^\bullet_{\text{poly}}\to\mathcal{D}^\bullet_{\text{poly}}$.

The general case is analogous. The statement of Lemma is true for any L_∞-morphism $\mathcal{U}\colon T^\bullet_{\text{poly}}\to\mathcal{D}^\bullet_{\text{poly}}$, not only for the Kontsevich's one. □

1.6.1. Now we have two modules over $T^\bullet_{\text{poly}}(\mathbb{R}^d)$: the usual module $\Omega^\bullet(\mathbb{R}^d)$, and the L_∞-module $C_\bullet(A,A)$. We want to construct an L_∞-morphism

$$\hat{\mathcal{U}}\colon C_\bullet(A,A)\to\Omega^\bullet(\mathbb{R}^d)$$

between them.

It means that we search for maps

$$\hat{\mathcal{U}}_k\colon\Lambda^k T^\bullet_{\text{poly}}(\mathbb{R}^d)\otimes C_\bullet(A,A)\to\Omega^\bullet(\mathbb{R}^d)[-k],\quad k\ge 0$$

such that for $k \geq -2$

(21)
$$\sum \pm\hat{\mathcal{U}}_{k+1}([\gamma_{i_1},\gamma_{i_2}]\wedge\gamma_{j_1}\wedge\cdots\wedge\gamma_{j_k}\otimes\omega)+\sum_{\substack{p+q=k+2,\\ q\geq 0}}\pm\hat{\mathcal{U}}_{p+1}(\gamma_{i_1}\wedge\cdots\wedge\gamma_{i_p}\otimes L_{\mathcal{U}_q(\gamma_{j_1}\wedge\cdots\wedge\gamma_{j_q})}\omega)+$$
$$+\sum\pm L_{\gamma_i}(\hat{\mathcal{U}}_{k+1}(\gamma_{j_1}\wedge\cdots\wedge\gamma_{j_{k+1}}\otimes\omega))=0 \quad (L_{\mathcal{U}_0}=b).$$

We set:

$$\hat{\mathcal{U}}_0(\omega)=\mu\omega \quad \text{(see Theorem 1.3.1)}.$$

For $k=-2$ (21) is just the statement that μ is a map of the complexes.

In the rest of this paper we construct such an L_∞-morphism $\hat{\mathcal{U}}$, where $\mathcal{U}$ is the Kontsevich formality morphism.

2. Construction of the morphism $\hat{\mathcal{U}}$

2.1. **The Kontsevich formality morphism $\mathcal{U}$.** Here we recall, very briefly, the construction [K] of the Kontsevich formality morphism $\mathcal{U}$.

We need to construct maps

$$\mathcal{U}_k\colon \Lambda^k T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)\to\mathcal{D}^\bullet_{\mathrm{poly}}(\mathbb{R}^d)[1-k].$$

The formula for $\mathcal{U}_k$ is organized as a sum over the admissible graphs Γ. We cite from [K] the definition of an admissible graph Γ.

2.1.1.

Definition. Admissible graph Γ is an oriented graph with labels such that

1) the set of vertices V_Γ is $\{1,\dots,n\}\sqcup\{\bar{1},\dots,\bar{m}\}$, $n,m\in\mathbb{Z}_{\geq 0}$, $2n+2+m\geq 0$; vertices from the set $\{1,\dots,n\}$ are called vertices of the first type, vertices from $\{\bar{1},\dots,\bar{m}\}$ are called vertices of the second type,
2) every edge $(v_1,v_2)\in E_\Gamma$ starts at a vertex of the first type,
3) there are no loops, i.e. no edges of the type (v,v),
4) for every vertex $k\in\{1,\dots,n\}$ of the first type, the set of edges
$$\mathrm{Star}(k)=\big\{(v_1,v_2)\in E_\Gamma \mid v_1=k\big\}$$
starting from k, is labeled by symbols $(e_k^1,\dots,e_k^{\#\mathrm{Star}(k)})$.

2.1.2. For any admissible graph Γ with n vertices of the first type and m vertices of the second type we define a map $\mathcal{U}_\Gamma\colon\gamma_1\otimes\gamma_2\otimes\cdots\otimes\gamma_n\mapsto\{A^{\otimes m}\to A\}$, $A=C^\infty(\mathbb{R}^d)$, and γ_i is $(\#\mathrm{Star}(i))$-polyvector field. The function $\Phi=\mathcal{U}_\Gamma(\gamma_1\otimes\cdots\otimes\gamma_n)(f_1\otimes\cdots\otimes f_m)$ is defined as follows. It is the sum over all configurations of indices running from 1 to d, labeled by E_Γ:

$$\Phi=\sum_{I\colon E_\Gamma\to\{1,\dots,d\}}\Phi_I$$

where Φ_I is the product over all $n+m$ vertices of Γ of certain partial derivatives of functions f_i and of coefficients of γ_i.

Namely, with each vertex i, $1\leq i\leq n$ of the first type we associate function Ψ_i an $\mathbb{R}^d$ which is

$$\Psi_i=\langle\gamma_i,dx^{I(e_i^1)}\otimes\cdots\otimes dx^{I(e_i^{\#\mathrm{Star}(i)})}\rangle.$$

For each vertex $\bar{j}$ of the second type the associated function $\Psi_{\bar{j}}$ is defined as f_j.

In the next step we put into each vertex v instead of function Ψ_v, its partial derivative

$$\tilde{\Psi}_v = \left(\prod_{e \in E_\Gamma,\ e=(*,v)} \partial_{I(e)} \right) \Psi_v.$$

Then, $\Phi_I = \prod_{v \in V_\Gamma} \tilde{\Psi}_v$, and $\Phi = \sum_I \Phi_I$.

This is just a usual construction of GL_d-invariants from graphs.

2.1.3. Till now, the number of edges of Γ with n vertices of the first type and m vertices of the second type was not fixed. We claim, that it is uniquely defined by the request that $\mathcal{U}_\Gamma$ is a map

$$\mathcal{U}_\Gamma \colon \Lambda^n T^\bullet_{\text{poly}}(\mathbb{R}^d) \to \mathcal{D}^\bullet_{\text{poly}}(\mathbb{R}^d)[1-n],$$

i.e. by the grading.

As an element of $\mathcal{D}^\bullet_{\text{poly}}(\mathbb{R}^d)$, $\mathcal{U}_\Gamma$ has grading $m-1$. This grading should be equal to $\deg \gamma_1 + \cdots + \deg \gamma_n + 1 - n$. We have:

$$\deg \gamma_i = \#\text{Star}(i) - 1,$$

and

$$\# E_\Gamma = \sum_{i \in \{1,\dots,n\}} \#\text{Star}(i),$$

because any edge starts at a vertex of the first type by Definition 2.1.1.

Therefore,

$$\# E_\Gamma = \deg \gamma_1 + \cdots + \deg \gamma_n + n.$$

But

$$m - 1 = \deg \gamma_1 + \cdots + \deg \gamma_n + 1 - n,$$

therefore

$$m - 1 = \# E_\Gamma - n + 1 - n,$$

and

$$\# E_\Gamma = 2n + m - 2.$$

2.1.4. Now we search a formula for $\mathcal{U}_n$ in the form

$$\mathcal{U}_n = \sum_\Gamma W_\Gamma \times \mathcal{U}_\Gamma, \tag{22}$$

where Γ has n vertices of the first type, and $W_\Gamma \in \mathbb{C}$ is a number. We want to define W_Γ as an integral of a form Ω_Γ of the top degree over a configuration space. Any edge of Γ will define a 1-form on this configuration space, and the form Ω_Γ is a wedge product of these 1-forms (in order corresponded to the labeling of the graph). Therefore, the number $\# E_\Gamma$ of the edges should be equal to dimension of the configuration space. Therefore, the configuration space should be a quotient by a 2-dimensional group of n "2-dimensional points" and m "1-dimensional" points.

2.1.5. Here we define these configuration spaces. Let $\mathcal{H}$ be the open complex upper half-plane.

First, denote by $\mathrm{Conf}_{n,m}$ the space

$$\begin{aligned}\mathrm{Conf}_{n,m} = \big\{p_1,\dots,p_n;q_1,\dots,q_m \mid p_i\in\mathcal{H} \text{ for all } i=1,\dots,n,\\ q_j\in\mathbb{R}=\partial\bar{\mathcal{H}} \text{ for all } j=1,\dots,m;\ p_{i_1}\ne p_{i_2} \text{ for } i_1\ne i_2,\ q_{j_1}\ne q_{j_2} \text{ for } j_1\ne j_2\big\},\end{aligned}$$

and

$$\mathrm{Conf}_n = \big\{p_1,\dots,p_n \mid p_i\in\mathbb{C}, p_{i_1}\ne p_{i_2} \text{ for } i_1\ne i_2\big\}.$$

For $2n+m\ge 2$ the group

$$G^{(1)} = \big\{z\mapsto az+b \mid a\in\mathbb{R}_{>0},\ b\in\mathbb{R}\big\}$$

acts freely on the space $\mathrm{Conf}_{n,m}$; therefore, the quotient-space

$$C_{n,m} = \mathrm{Conf}_{n,m}/G^{(1)}$$

is a smooth manifold of dimension $2n+m-2$. Analogously,

$$C_n = \mathrm{Conf}_n/G^{(2)}$$

where

$$G^{(2)} = \big\{z\mapsto az+b \mid a\in\mathbb{R}_{>0},\ b\in\mathbb{C}\big\}.$$

It is a smooth manifold for $n\ge 2$.

M. Kontsevich constructed in [K], Section 6, a compactification of these spaces. The differential forms, defined below, can be prolonged to regular differential forms on these compactifications. Therefore, the integrals (defined below) converge.

We will not discuss these compactifications here. We will consider this question for other spaces, suitable for the construction of map $\hat{\mathcal{U}}$, in Section 2.2.

2.1.6. Now, for a graph Γ with n vertices of the first type and m vertices of the second type, we are going to define a form Ω_Γ of the top degree on $C_{n,m}$, and then set $W_\Gamma = \sim\int_{C_{n,m}}\Omega_\Gamma$ (up to some coefficient which depends on n,m and does not depend on Γ).

For any two points $p,q\in\bar{\mathcal{H}}$, denote by $l(p,q)$ the geodesic in the Poincaré metric on $\mathcal{H}$, passing through the points p and q. It is just a (part of) half-circle, passing through p and q, and orthogonal to $\mathbb{R}=\partial\bar{\mathcal{H}}$. Denote by $l(p,\infty)$ the vertical line passing through the point p. Denote by $\phi^h(p,q)\in\mathbb{R}/2\pi\mathbb{Z}$ the angle between $l(p,q)$ and $l(p,\infty)$, from $l(p,\infty)$ to $l(p,q)$ counterclockwise. The formula for $\phi^h(p,q)$ is

$$\phi^h(p,q) = \frac{1}{2i}\,\mathrm{Log}\left(\frac{(q-p)(\bar{q}-p)}{(q-\bar{p})(\bar{q}-\bar{p})}\right). \tag{23}$$

The 1-form $d\phi^h(p,q)$ is well-defined.

Now we set that the vertices $\{1,\dots,n\}$ of the first type of the graph Γ are the points $p_1,\dots,p_n$ on $\mathrm{Conf}_{n,m}$, and vertices $\{\bar{1},\dots,\bar{m}\}$ are the points $q_1,\dots,q_m$ on $\mathrm{Conf}_{n,m}$. In other words, the graph "is placed" on $\mathcal{H}\sqcup\mathbb{R}$ in all possible ways. Then each edge $e=(p,q)$ of Γ defines a map from $\mathrm{Conf}_{n,m}$ to $\mathrm{Conf}_{2,0}$ or to $\mathrm{Conf}_{1,1}$. On both spaces $\mathrm{Conf}_{2,0}$ and $\mathrm{Conf}_{1,1}$ we have constructed the 1-form ϕ^h. Denote by φ_e its pull-back to $\mathrm{Conf}_{n,m}$. It is clear that φ_e is $G^{(1)}$-invariant because $G^{(1)}$ is the group of transformations in the Poincaré metric on $\mathcal{H}$ preserving $\{\infty\}$. Thus, φ_e is a well-defined 1-form on $C_{n,m}=\mathrm{Conf}_{n,m}/G^{(1)}$.

We set:

$$W_\Gamma = \prod_{k=1}^{n} \frac{1}{(\#\mathrm{Star}(k))!} \cdot \frac{1}{(2\pi)^{2n+m-2}} \cdot \int_{C^+_{n,m}} \bigwedge_{e\in E_\Gamma} \varphi_e. \tag{24}$$

Here $C^+_{n,m}$ is the connected component of $C_{n,m}$ where $q_1 < \cdots < q_m$.

Theorem (M. Kontsevich,[K]). *The maps $\mathcal{U}_k$, $k \geq 1$, defined by formulas* (22), (24), *are the Taylor components of an L_∞-morphism $\mathcal{U}\colon T^\bullet_{\mathrm{poly}}(\mathbb{R}^d) \to \mathcal{D}^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$. That is they satisfy the identity* (16).

2.1.7. One of the crucial points is that $G^{(1)}$ is the full group of symmetries for the 1-form $d\phi^h$. This is why the integrals (24) a priori do not vanish (and they actually do not).

2.2. Construction of the map $\hat{\mathcal{U}}$.

2.2.1. *Configuration spaces $D_{\mathbf{1},n,m}$.* Let D^2 be the open disk $D^2 = \{z \in \mathbb{C} \mid |z| < 1\}$. We denote by $\mathbf{1}$ the center of this disk $\mathbf{1} = \{z = 0\}$. (It's a notation bad from any point of view). The space $\mathrm{Disk}_{\mathbf{1},n,m}$ is

$$\mathrm{Disk}_{\mathbf{1},n,m} = \{p_1,\dots,p_n \in D^2,\ q_1,\dots,q_m \in S^1 = \partial\bar{D}^2,$$
$$p_i \neq p_j \text{ for } i\neq j,\ p_i \neq \mathbf{1} \text{ for any } i,\ q_i \neq q_j \text{ for } i \neq j\}.$$

For $2n+m \geq 1$ the group of rotations

$$G = \{z \mapsto e^{i\theta} z,\ \theta \in \mathbb{R}/2\pi\mathbb{Z}\}$$

acts freely on $\mathrm{Disk}_{\mathbf{1},n,m}$, and we define the manifold $D_{\mathbf{1},n,m}$ as

$$D_{\mathbf{1},n,m} = \mathrm{Disk}_{\mathbf{1},n,m}/G. \tag{25}$$

It has dimension $2n+m-1$. Note that $G\mathbf{1} = \mathbf{1}$.

Now we are going to describe a compactification $\bar{\mathcal{D}}_{\mathbf{1},n,m}$ which will be used in the sequel.

The idea arise from [K], Section 5; we just describe all strata of codimension 1.

S1) some points $p_{i_1},\dots,p_{i_k}$ of the first type move close to each other and far from $\mathbf{1}$, $k \geq 2$. The corresponding boundary stratum is $C_k \times D_{\mathbf{1},n-k+1,m}$;

S2) some points $p_{i_1},\dots,p_{i_k}$ of the first type, $k\geq 1$, move close to each other and close to $\mathbf{1}$. To describe this stratum denote by D_k the following space:

$$D_k = \{p_1,\dots,p_k \in \mathbb{C},\ p_i \neq p_j \text{ for } i \neq j\}/\{z \mapsto az,\ a \in \mathbb{R}_{>0}\}.$$

For $k \geq 1$ it is a manifold of dimension $k-1$. The boundary stratum in the case S2) is $D_k \times D_{\mathbf{1},n-k,m}$;

S3) some points $p_{i_1},\dots,p_{i_k}$ of the first type and some points $q_{j_1},\dots,q_{j_l}$ of the second type move close to each other (and close to $S^1 = \partial\bar{D}^2$), $2k+l \geq 2$. In this case the boundary stratum is

$$C_{k,l} \times D_{\mathbf{1},n-k,m-l+1}.$$

One can check that in all three cases the described strata have codimension 1.

In the general case of arbitrary codimension, we have several groups of points moving close to each other; this gives strata of codimension > 1. Then "we are looking through a magnifying glass" to each group and find that inside it there are several groups of points, moving close to each other, an so on. All the boundary strata can be described by labeled and colored trees.

For us it is principal that the differential 1-forms on $D_{\mathbf{1},n,m}$ (analogs of $d\phi^h$) constructed below can be prolonged to regular differential 1-forms on the compactification $\bar{D}_{\mathbf{1},n,m}$. The proof will be clear after the definition of these 1-forms.

2.2.2. *Admissible graphs.* An admissible graph is the same that in [K] (see Section 2.1.1) but here we have a marked vertex $\mathbf{1}$, and $\mathbf{1}$ is not the end-point for any edge. There are vertex $\mathbf{1}$, vertices of the first type $\{1,\dots,n\}$, and vertices of the second type $\{\bar{1},\dots,\bar{m}\}$.

2.2.3. *Differential forms.* There are two types of edges: connecting $\mathbf{1} \in D^2$ with an other point, and all others edges. We will define a differential 1-form φ_e separately in these two cases.

C1) e is an usual edge, i.e.

$$e = (p,q),\ \ p,q \in \bar{D}^2,\ \ p \neq \mathbf{1}.$$



Figure 1.

In this case we consider the angle θ_e between the geodesic $l(p,q)$ in the Poincaré metric on D^2, with the geodesic $l(p,\mathbf{1})$ (the last geodesic should be a diametral line), counted from $l(p,\mathbf{1})$ to $l(p,q)$ counterclockwise. The angle θ_e is defined modulo 2π. By definition, $\varphi_e = d\theta_e$; it is a well-defined 1-form on the space $D_{\mathbf{1},2,0}$, or on $D_{\mathbf{1},1,1}$, see Figure 1.

C2) $e = (p,q)$, $p = \mathbf{1}$.



FIGURE 2.

In this case θ_e is the angle between the line $(\mathbf{1}, q)$ and the line $(\mathbf{1}, a_1)$ where $a_1 = \{\bar{1}\}$, the first vertex of the second type on S^1. The 1-form $\varphi_e = d\theta_e$ is well-defined 1-form on $D_{\mathbf{1},1,1}$ or $D_{\mathbf{1},0,2}$. We denote by φ_e also its pull-back on $D_{\mathbf{1},n,m}$, see Figure 2.

2.2.4. *Polydifferential operators, corresponded to the graphs.* Now we want to define some operators

$$\Lambda^n T^{\bullet}_{\mathrm{poly}}(\mathbb{R}^d) \otimes C_{\bullet}(A, A) \to \Omega^{\bullet}(\mathbb{R}^d)[-n].$$

Let $\gamma_1, \dots, \gamma_n$ be polyvector fields on $\mathbb{R}^d$, and let $a_1, \dots, a_m$ be m functions, $a_i \in A = C^\infty(\mathbb{R}^d)$. Now let Γ be an admissible graph, such that $\#\mathrm{Star}(i) = \deg \gamma_i + 1$ for a vertex i of the first type, $\#\mathrm{Star}(\mathbf{1}) = l$. Then we are going to construct an outcoming l-differential form Ω^Γ_l. To do that, we define $\Omega^\Gamma_l(\partial_{\alpha_1} \wedge \dots \wedge \partial_{\alpha_l})$, $i_1, \dots, i_l \in \{1, \dots, d\}$. As in Section 2.1.2,

$$\Omega^\Gamma_l(\partial_{\alpha_1} \wedge \dots \wedge \partial_{\alpha_l}) = \sum_{I\colon E_\Gamma \setminus \mathrm{Star}(\mathbf{1}) \to \{1,\dots,d\}} \Omega_I^{\alpha_1,\dots,\alpha_l}$$

We can extend the map $I\colon E_\Gamma \setminus \mathrm{Star}(\mathbf{1}) \to \{1, \dots, d\}$ to a map $\hat{I}\colon E_\Gamma \to \{1, \dots, d\}$, where if $e = (\mathbf{1}, *)$ and e has the label $e^s_{\mathbf{1}}$ in the graph Γ, we set $I(e) = \alpha_s$. Then the functions Ψ_v, $v \neq \mathbf{1}$, are defined as in Section 2.1.2. Next,

$$\tilde{\Psi}_v = \left(\prod_{e \in E_\Gamma,\ e=(*,v)} \partial_{\hat{I}(e)} \right) \Psi_v, \quad v \neq \mathbf{1},$$

and then

$$\Omega_I^{\alpha_1,\dots,\alpha_l} = \prod_{v \in V_\Gamma \setminus \mathbf{1}} \tilde{\Psi}_v.$$

Finally,

$$\Omega_l^\Gamma(\partial_{\alpha_1}\wedge\dots\wedge\partial_{\alpha_l})=\sum_I \Omega_I^{\alpha_1,\dots,\alpha_l}.$$

2.2.5. *The grading.* Let us find a relation on $n, m, \#E_\Gamma$ and l. We want to define a map

$$\Lambda^n T^\bullet_{\mathrm{poly}}\otimes C_{m-1}(A,A)\to\Omega^\bullet[-n].$$

Therefore,

$$\deg\gamma_1+\dots+\deg\gamma_n+(-m+1)-n=-l. \tag{26}$$

On the other hand

$$\#E_\Gamma=\deg\gamma_1+\dots+\deg\gamma_n+n+l. \tag{27}$$

We have from (26) and (27):

$$\#E_\Gamma-n-l-m+1-n=-l \tag{28}$$

or

$$\#E_\Gamma=2n+m-1. \tag{29}$$

Therefore, for the grading condition a graph with n vertices of the first type and m vertices of the second type should have $2n+m-1$ edges. According to our discussion in Section 2.1.4, the configuration space for the construction of $\hat{\mathcal{U}}$ should have dimension $2n+m-1$. The space $D_{\mathbf{1},n,m}$ has precisely this dimension.

2.2.6. *Weight W_Γ.* Now we consider a graph Γ as placed on the space $\mathrm{Disk}_{\mathbf{1},n,m}$ such that the vertex $\mathbf{1}$ of Γ is the center $\mathbf{1}$ of the disk D^2, the points $\{p_1,\dots,p_n\}$ of $\mathrm{Disk}_{\mathbf{1},n,m}$ are the vertices $\{1,\dots,n\}$ of Γ of the first type, and the points $\{q_1,\dots,q_m\}$ of $\mathrm{Disk}_{\mathbf{1},n,m}$ are the vertices $\{\bar{1},\dots,\bar{m}\}$ of Γ of the second type. Then each edge e of Γ defines a map

$$pr_e\colon \mathrm{Disk}_{\mathbf{1},n,m}\to\mathrm{Disk}_{\mathbf{1},2,0}\quad\text{or}$$
$$pr_e\colon \mathrm{Disk}_{\mathbf{1},n,m}\to\mathrm{Disk}_{\mathbf{1},1,1}$$

We denote by the same symbol φ_e the pull-back $\mathrm{pr}_e^*\varphi_e$ of the 1-form, constructed in Section 2.2.3. The 1-form φ_e is G-invariant (G is the rotation group), and we can consider φ_e as a 1-form on the space $\mathcal{D}_{\mathbf{1},n,m}$.

Definition (the weight W_Γ)**.**

$$W_\Gamma=\frac{1}{(\#\mathrm{Star}(\mathbf{1}))!}\prod_{k=1}^n\frac{1}{(\#\mathrm{Star}(k))!}\cdot\frac{1}{(2\pi)^{2n+m-1}}\cdot\int_{D^+_{\mathbf{1},n,m}}\bigwedge_{e\in E_\Gamma}\varphi_e \tag{30}$$

where $D^+_{\mathbf{1},n,m}$ is the connected component of $D_{\mathbf{1},n,m}$ for which the points $(q_1,\dots,q_m)\in S^1$ define the right cyclic order on S^1.

2.2.7.

Theorem. *Let $G_{\mathbf{1},n,m}$ be the set of the admissible graphs in the sence of Section* 2.2.2 *with $2n+m-1$ edges. Define the map*

$$\hat{\mathcal{U}}_n\colon \Lambda^n T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)\otimes C_{m-1}(A,A)\to \Omega^\bullet(\mathbb{R}^d)[-n]$$

by the formula

$$\hat{\mathcal{U}}_n=\sum_{\Gamma\in G_{\mathbf{1},n,m}} W_\Gamma\times\Omega_l^\Gamma, \tag{31}$$

$$l=n+m-1-\sum_{i=1}^n \deg\gamma_i. \tag{32}$$

Then the maps $\{\hat{\mathcal{U}}_n\}$ are the Taylor components of an L_∞-morphism of the L_∞-modules over $T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$:

$$\hat{\mathcal{U}}\colon C_\bullet(A,A)\to\Omega^\bullet(\mathbb{R}^d).$$

The L_∞-module over $T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ structure on $C_\bullet(A,A)$ is defined by (19) *where $\{\mathcal{U}_k\}$ are the Taylor components of the Kontsevich formality morphism* (*with the harmonic angle function, as in Section* 2.1).

Proof. We need to prove (21) for $\{\hat{U}_n\}$ defined as above. The l.h.s. of (21) has a form

$$\sum_{\Gamma\in G'_{\mathbf{1},n,m}} C_\Gamma\cdot\Omega_l^\Gamma(\gamma_1\wedge\dots\wedge\gamma_n\otimes(a_1\otimes\dots\otimes a_m))$$

where $G'_{\mathbf{1},n,m}$ is the set of admissible graphs with n vertices of the first type, m vertices of the second type, and $2n+m-2$ edges, and $l=n+m-2-\sum_{i=1}^n\deg\gamma_i$. The numbers C_Γ are linear-quadratic expressions in the weights W_{Γ_1},defined in Section 2.2.6, and the Kontsevich weights W_{Γ_2}, defined in Section 2.1.6. We want to prove that $C_\Gamma=0$ for any $\Gamma\in G'_{\mathbf{1},n,m}$. The idea arise to [K]: we identify C_Γ with the integral over the boundary $\partial\bar{D}_{\mathbf{1},n,m}$ of the differential form of degree $2n+m-2$, namely, with $\bigwedge_{e\in E_\Gamma} d\varphi_e$. We have by the Stokes formula:

$$\int_{\partial\bar{D}_{\mathbf{1},n,m}}\bigwedge_{e\in E_\Gamma} d\varphi_e=\int_{\bar{D}_{\mathbf{1},n,m}} d\left(\bigwedge_{e\in E_\Gamma} d\varphi_e\right)=0. \tag{33}$$

On the other hand, only the boundary strata of codimension 1 contributes in the l.h.s. of (33). We have:

(34)

$$\int_{\partial\bar{D}_{\mathbf{1},n,m}}\bigwedge_{e\in E_\Gamma} d\varphi_e=\int_{\partial_{\mathrm{S1)}}\bar{D}_{\mathbf{1},n,m}}\bigwedge_{e\in E_\Gamma} d\varphi_e+\int_{\partial_{\mathrm{S2)}}\bar{D}_{\mathbf{1},n,m}}\bigwedge_{e\in E_\Gamma} d\varphi_e+\int_{\partial_{\mathrm{S3)}}\bar{D}_{\mathbf{1},n,m}}\bigwedge_{e\in E_\Gamma} d\varphi_e$$

where S1), S2), and S3) are the three types of the boundary strata of codimension 1, listed in Section 2.2.1. The idea of the proof is to identify these summands with the summands contributed to C_Γ from (21). We consider the cases S1–S3) separately.

2.2.7.1. **The case S1).** The boundary stratum is $C_k\times D_{\mathbf{1},n-k+1,m}$. It is clear that the integral over it factorizes in the product of two integrals: the integral over C_k and the integral over $D_{\mathbf{1},n-k+1,m}$. It is proved in [K], Section 6, that the integral over C_k

does not vanish only for $k = 2$. The situation is as follows: two points move close to each other and far from $\mathbf{1}$. This corresponds to the first summand of (21), containing the Schouten–Nijenhuis bracket of polyvector fields.

2.2.7.2. **The case S2).** Some points $p_{i_1}, \dots, p_{i_k}$ move close to each other and close to $\mathbf{1} \in D^2$. The boundary stratum is $D_k \times D_{\mathbf{1},n-k,m}$. It is clear that the integral factorizes into the product of two integrals, of an integral over D_k and of an integral over $D_{\mathbf{1},n-k,m}$ (see, in particular, the following two subsections). Then, by Theorem 6.6.1 in [K], we have only the following possibilities:

S2.1) $k = 1$ and there is an edge from $\mathbf{1}$ to p_{i_1};
S2.2) $k = 1$ and there are no edges connecting p_{i_1} and $\mathbf{1}$.

We analyze these strata separately below. It will be clear from our discussion how to apply Theorem 6.6.1 in [K] to prove that other possible strata give zero contribution.

2.2.7.2.1 **The case S2.1).**



Figure 3.

Let (p_{i_1}, q) be an edge starting at p_{i_1}, and let Φ_1 be the angle between $(\mathbf{1}, p_{i_1})$ and $(\mathbf{1}, a_1)$ (this is the angle θ_e for $e = (\mathbf{1}, p_{i_1})$), Φ_2 be the angle between $(\mathbf{1}, q)$ and $(\mathbf{1}, a_1)$, and Φ_3 be the angle between (p_{i_1}, q) and $(p_{i_1}, \mathbf{1})$, see Figure 3. The integral over D_1

should be the integral over D_1 of the canonical 1-form $d\varphi$ on D_1. By our definition, θ_e for $e = (p_{i_1}, q)$ is the angle Φ_3. The 2-form $d\varphi_{(p_{i_1},q)} \wedge d\varphi_{(\mathbf{1},p_{i_1})}$ is equal to $d\Phi_3 \wedge d\Phi_1$. In the limit $p_{i_1} \to \mathbf{1}$ the angle $\Phi_3 \to \pi - (\Phi_1 - \Phi_2)$ (see Fig. 3). Therefore, we have two-form $d(\pi - \Phi_1 + \Phi_2) \wedge d\Phi_1 = d\Phi_2 \wedge d\Phi_1$. The angle Φ_2 does not depend on the position of p_{i_1}. The "new" edge will be $(\mathbf{1}, q)$, and its angle is precisely Φ_2. The integral over D_1 is $\int d\Phi_1$.

In the general case, there are some edges starting at p, but no edge ends at p: in this case the angle of this edge would be equal to 0.

These terms correspond to a summand of $d \circ i_{\gamma_i}(\hat{\mathcal{U}}_{k+1}(\gamma_{j_1} \wedge \dots \wedge \gamma_{j_{k+1}} \otimes \omega))$ (see (21)) and any they contribute to the third summand in (21). See Section 2.2.7.2.3 for a further discussion.

2.2.7.2.2 **The case S2.2).** We denote p_{i_1} by p. There no edges connecting p and $\mathbf{1}$. See Figure 4.



FIGURE 4.

In the limit when p tends to $\mathbf{1}$ and the points $q_1, q_2, q_3 \dots$ (end-points of the edges starting from p) are far from $\mathbf{1}$, the wedge product $d\phi_1 \wedge d\phi_2 \wedge d\phi_3 \wedge \dots$ tends to $\pm d\phi_0 \wedge d(\phi_2 - \phi_1) \wedge d(\phi_3 - \phi_1)$. The integral $\int_{S^1} d\phi_0$ is the integral over D_1. The remaining part of the wedge product, $d(\phi_2-\phi_1)\wedge d(\phi_3-\phi_1)\wedge\dots$ contributes to the integral over $D_{\mathbf{1},n-1,m}$.

In the general case, if there exists any edge ending at p, an edge $l=(s,p)$, and s is far from $\mathbf{1}$, the angle φ_e tends to 0 when p tends to $\mathbf{1}$. Therefore, no edge ends at p.

These terms correspond to a summand of $i_{\gamma_i}\circ d(\hat{\mathcal{U}}_{k+1}(\gamma_{j_1}\wedge\dots\wedge\gamma_{k+1}\otimes\omega))$ and contribute to the third summand in (21). See Section 2.2.7.2.3 for a further discussion.

2.2.7.2.3 We claim that the total contribution of the two cases S2.1) and S2.2) is equal exactly to $L_{\gamma_i}\tilde{\omega}=d\circ i_{\gamma_i}\tilde{\omega}\pm i_{\gamma_i}\circ d\tilde{\omega}$, where $\tilde{\omega}=\hat{\mathcal{U}}_{k+1}(\gamma_{j_1}\wedge\dots\wedge\gamma_{j_{k+1}}\otimes\omega)$. Here we consider two examples.

Example 1. $\gamma_i=v$ is a vector field, $\tilde{\omega}$ is a function. Then $d\circ i_v\tilde{\omega}=0$, and we have only the summand $i_v\circ d\tilde{\omega}$, which corresponds exactly exactly to the case S2.2).

Example 2. Again, $\gamma_i=v$ is a vector field, $v=\sum_i v_i\partial_i$, but $\tilde{\omega}$ is a 1-form $\tilde{\omega}=\sum_i f_i dx_i$. Then

$$(*)\qquad d\circ i_v\tilde{\omega}=d\left(\sum_i v_if_i\right)=\sum_{i,j}f_i\frac{\partial v_i}{\partial x_j}dx_i+\sum_{i,j}v_i\frac{\partial f_i}{\partial x_j}dx_j$$

On the other hand,

$$(**)\qquad i_vd\omega=i_v\left(\sum_{i,j}\frac{\partial f_i}{\partial x_j}dx_j\wedge dx_i\right)=\sum_{i,j}\frac{\partial f_i}{\partial x_j}\cdot v_jdx_i-\sum_{i,j}\frac{\partial f_i}{\partial x_j}v_idx_j$$

We see that in $L_v\tilde{\omega}=d\circ i_v\tilde{\omega}+i_v\circ d\tilde{\omega}$ the second summand in the r.h.s. of $(*)$ cancells the second summand in the r.h.s of $(**)$, whereas the first summand in the r.h.s. of $(*)$ corresponds to the case S2.1), and the first summand in the r.h.s. of $(**)$ corresponds to the case S2.2).

In the general case the situation is analogous to this example.

2.2.7.3. **The case S3).** Some points $p_{i_1},\dots,p_{i_k}$ of the first type and some points $q_{j_1},\dots,q_{j_l}$ of the second type move close to each other and close to $S^1=\partial\bar{D}^2$, $2k+l\geq 2$. The boundary stratum is $C_{k,l}\times D_{\mathbf{1},n-k,m-l+1}$. The integral over $C_{k,l}$ is exactly the Kontsevich integral: the angle with $\mathbf{1}$ becomes the angle with ∞ for $C_{k,l}$. We obtain the second summand in (21).

Theorem 2.2.7 is proven. □

2.2.8. It is clear that

$$\hat{\mathcal{U}}_0(a_1\otimes\dots\otimes a_m)=\mu(a_1\otimes\dots\otimes a_m)=\frac{1}{(m-1)!}a_1\,da_2\wedge\dots\wedge da_m.$$

3. APPLICATIONS TO TRACES, CUP-PRODUCTS, AND TO THE DUFLO FORMULA

3.1. First of all, we recall what the (extended) Duflo formula is. Let $\mathfrak{g}$ be a finite-dimensional Lie algebra, we associate with $\mathfrak{g}$ two $\mathfrak{g}$-modules: the symmetric algebra $S^\bullet(\mathfrak{g})$, and the universal enveloping algebra $U(\mathfrak{g})$. These two $\mathfrak{g}$-modules are isomorphic due to the Poincaré–Birkhoff–Witt theorem, the isomorphism $\varphi_{PBW}\colon S^\bullet(\mathfrak{g})\to U(\mathfrak{g})$ is defined as

$$\varphi_{PBW}(g_1\cdot\dots\cdot g_k)=\frac{1}{k!}\sum_{\sigma\in\Sigma_k}g_{\sigma(1)}\otimes\dots\otimes g_{\sigma(k)}.$$

As algebras, $S^\bullet(\mathfrak{g})$ and $U(\mathfrak{g})$ are not isomorphic. The Duflo theorem states that the algebras of $\mathfrak{g}$-invariants $[S^\bullet(\mathfrak{g})]^\mathfrak{g}$ and $[U(\mathfrak{g})]^\mathfrak{g}$ are (canonically) isomorphic. Let us recall the construction of this isomorphism.

For any $k \geq 1$, let Tr_k be a canonical invariant element $\mathrm{Tr}_k \in [S^k(g^*)]^\mathfrak{g}$, defined as the symmetrization of the map $g \mapsto \mathrm{Tr}|_\mathfrak{g} \, \mathrm{ad}^k g$ $(g \in \mathfrak{g})$. We consider elements of $S^k(\mathfrak{g}^*)$ as differential operators of k-th order on $S^\bullet(\mathfrak{g})$ with constant coefficients. We define the map $\varphi_{\text{strange}} \colon S^\bullet(\mathfrak{g}) \to S^\bullet(\mathfrak{g})$ as

$$\varphi_{\text{strange}} = \exp\left(\sum_{k\geq 1} \alpha_{2k} \mathrm{Tr}_{2k}\right)$$

where the rational numbers α_{2k} are defined as

$$\sum_{k\geq 1} \alpha_{2k} x^{2k} = \frac{1}{2} \log \frac{e^{\frac{x}{2}} - e^{-\frac{x}{2}}}{x}.$$

It is clear that $\varphi_{\text{strange}} \colon S^\bullet(\mathfrak{g}) \to S^\bullet(\mathfrak{g})$ is a map of $\mathfrak{g}$-modules, because $\mathrm{Tr}_{2k} \in [S^{2k}(\mathfrak{g}^*)]^\mathfrak{g}$.

3.1.1.

Theorem (Duflo)**.** *For any finite-dimensional Lie algebra* $\mathfrak{g}$ *the map*

$$\varphi_D^* = [\varphi_{PBW} \circ \varphi_{\text{strange}}] \colon [S^\bullet(\mathfrak{g})]^\mathfrak{g} \to [U(\mathfrak{g})]^\mathfrak{g}$$

is an isomorphism of algebras.

3.1.2.

Theorem (Kontsevich [K])**.** *For any finite-dimensional Lie algebra* $\mathfrak{g}$ *the map* φ_D *of* $\mathfrak{g}$*-modules* $\varphi_D \colon S^\bullet(\mathfrak{g}) \to U(\mathfrak{g})$ *defines the map of algebras*

$$\varphi_D^* \colon H^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g})) \to H^\bullet(\mathfrak{g}; U(\mathfrak{g})).$$

3.1.3. Now consider *homology* $H_\bullet(\mathfrak{g}; S(\mathfrak{g}))$ and $H_\bullet(\mathfrak{g}; U(\mathfrak{g}))$. They have a structure of modules over the algebras $H^\bullet(\mathfrak{g}, S(\mathfrak{g}))$ and $H^\bullet(\mathfrak{g}; U(\mathfrak{g}))$, correspondingly. In particular, for zero (co)homology $A^{\text{coinv}} = A/\mathfrak{g} \cdot A$ has a structure of a module over the algebra A^{inv} for any $\mathfrak{g}$-module A which is an associative algebra such that the multiplication $A \otimes A \to A$ is a map of $\mathfrak{g}$-modules.

We want to construct a map of modules over $H^\bullet(\mathfrak{g}, \dots)$

$$\varphi_{D*} \colon H_\bullet(\mathfrak{g}; S(\mathfrak{g})) \to H_\bullet(\mathfrak{g}; U(\mathfrak{g}))$$

such that it is compatible with the map of algebras

$$\varphi_D^* \colon H^\bullet(\mathfrak{g}; S(\mathfrak{g})) \to H^\bullet(\mathfrak{g}, U(\mathfrak{g})).$$

It means that for any $\omega \in H^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g}))$ and $\eta \in H_\bullet(\mathfrak{g}, S^\bullet(\mathfrak{g}))$ one has

$$\varphi_{D*}(\omega \cdot \eta) = \varphi_D^*(\omega) \cdot \varphi_{D*}(\eta). \tag{35}$$

3.1.4.

Conjecture. *For any finite-dimensional Lie algebra* $\mathfrak{g}$*, the map* φ_{D*}*, induced by the map* $\varphi_D = \varphi_{PBW} \circ \varphi_{\mathrm{strange}}$ *of* $\mathfrak{g}$*-modules, satisfies* (35).

This conjecture is a typical application of a conjecture on cup-products on tangent cohomology, Conjecture 3.5.3.1 below. For a semisimple (or, more generally, unimodular) Lie algebra $\mathfrak{g}$ Conjecture 3.1.4 follows from Theorem 3.1.2: for example, the top-cohomology $H^{top}(\mathfrak{g}; A) \simeq H_0(\mathfrak{g}; A)$, and this is is an isomorphism of $H^0(\mathfrak{g}; A)$-modules for any A with compatible structures of an associative algebra and a $\mathfrak{g}$-module. Analogously for higher (co)homology.

For a general Lie algebra $\mathfrak{g}$, $H^{top}(\mathfrak{g}; M) \simeq H_0(\mathfrak{g}; \mathrm{Tr} \otimes M)$ where $\mathrm{Tr}\colon \mathfrak{g} \to \mathbb{C}$ is a one-dimensional $\mathfrak{g}$-module, the trace of the adjoint action. Therefore, for a general Lie algebra $\mathfrak{g}$, Theorem 3.1.2 does not imply Conjecture 3.1.4.

3.2. **L_∞-quasi-isomorphisms of L_∞-modules, the tangent complexes, the tangent map.** We say that an L_∞-map of two L_∞-modules $M^\bullet, N^\bullet$ (see Section 1.5.4) is an *L_∞-quasi-isomorphism of modules* if $\varphi_0\colon M^\bullet \to N^\bullet$ (which a priori is a map of complexes) is a quasi-isomorphism of complexes.

Let $M^\bullet$ be an L_∞-module over a dg Lie algebra $\mathfrak{g}$, and let $\pi \in \mathfrak{g}^1$ satisfies the Maurer–Cartan equation $d\pi + \frac{1}{2}[\pi, \pi] = 0$. We define the *tangent complex* $T_\pi M^\bullet$ as the graded space $M^\bullet$ with the differential d_π defined as follows:

$$d_\pi(m) = \phi_0(m) + \phi_1(\pi, m) + \frac{1}{2}\phi_2(\pi, \pi, m) + \dots + \frac{1}{n!}\phi_n(\pi, \pi, \dots, \pi, m) + \dots \tag{36}$$

where $\phi_k\colon \Lambda^k\mathfrak{g} \otimes M^\bullet \to M^\bullet[1-k]$ are the Taylor components of the L_∞-module structure on $M^\bullet$ (see Section 1.5.3).

Lemma.

(i) $d_\pi^2 = 0$ *if* $d\pi + \frac{1}{2}[\pi, \pi] = 0$;

(ii) *the formula*

$$(T_\pi\varphi)(a) = \sum_{k=0}^{\infty} \varphi_{k+1}(\pi, \pi, \dots, \pi, a)/k! \tag{37}$$

defines a map of the tangent complexes

$$T_\pi\varphi\colon T_\pi M^\bullet \to T_\pi N^\bullet;$$

(iii) *let* φ *be an* L_∞*-quasi-isomorphism of modules. Then* $T_\pi\varphi$ *is a quasi-isomorphism of the complexes provided the condition that* π *is sufficiently small. In particular, if we replace* π *on* $\hbar\pi$*, where* $\hbar$ *is a formal parameter,* $T_{\hbar\pi}\varphi$ *is a quasi-isomorphism.*

Proof. It is straightforward. □

3.3. From now on, we will work with formal power series in a formal parameter $\hbar$.

Lemma.

(i) *Let* $A = C^\infty(\mathbb{R}^d)[[\hbar]]$, $\pi \in \Lambda^2 T_{\mathrm{poly}}(\mathbb{R}^d)$, $[\pi, \pi] = 0$. *Then the map*

$$T_{\hbar\pi}\hat{\mathcal{U}}\colon T_{\hbar\pi}C_\bullet(A, A) \to T_{\hbar\pi}\Omega^\bullet(\mathbb{R}^d)[[\hbar]]$$

is a quasi-isomorphism of the complexes;

(ii)
$$T_{\hbar\pi}C_\bullet(A,A) = C_\bullet(A_*,A_*)$$
where A_* is the deformed algebra A with the Kontsevich star-product;

(iii)
$$T_{\hbar\pi}\Omega^\bullet(\mathbb{R}^d)[[\hbar]] = (\Omega^\bullet(\mathbb{R}^d)[[\hbar]], d = l_{\hbar\pi}).$$

Proof. (i) follows from Lemma 3.1, because $\hat{\mathcal{U}}_0 = \mu$ is a quasi-isomorphism by Theorem 1.3.1.

(ii) and (iii) follow from the definitions. □

In this way, we obtained a quasi-isomorphism of complexes:
$$T_{\hbar\pi}\hat{\mathcal{U}}\colon C_\bullet(A_*,A_*) \to (\Omega^\bullet(\mathbb{R}^d)[[\hbar]], d = L_{\hbar\pi}).$$

3.4.

Lemma.

(i)
$$H_0(C_\bullet(A_*,A_*)) = A_*/[A_*,A_*];$$

(ii)
$$H_0(\Omega^\bullet(\mathbb{R}^d)[[\hbar]], d = L_{\hbar\pi}) = A/\{A,A\},$$
where $\{A,A\}$ is the commutant of the Poisson algebra.

Proof. (i) In $C_\bullet(A_*,A_*)$ we have:
$$d(a_0 \otimes a_1) = a_0 * a_1 - a_1 * a_0.$$

(ii) By definition, $\operatorname{Im}(d\colon \Omega^1 \to \Omega^0) = L_{\hbar\pi}(\Omega^1)$. We have:
$$L_{\hbar\pi}(f dg) = (d_{\mathrm{DR}} \circ i_{\hbar\pi} + i_{\hbar\pi} \circ d_{\mathrm{DR}})(f dg) = i_{\hbar\pi} \circ d_{\mathrm{DR}}(f dg) = \hbar\pi(df \wedge dg) = \hbar\{f,g\}.$$
□

3.4.1.

Theorem. *The map $T_{\hbar\pi}\hat{\mathcal{U}}$ gives an isomorphism $A_*/[A_*,A_*] \xrightarrow{\sim} A/\{A,A\}$ for any Poisson bivector π.* □

We return to this map for a linear Poisson structure π in Section 3.6 after a conjecture on the compatibility with cup-products on the level of cohomology.

3.5. Cup-products on the tangent cohomology.

3.5.1.

Definition. For a dg Lie algebra $\mathfrak{g}$, and a solution π of the Maurer–Cartan equation $d\pi + \frac{1}{2}[\pi,\pi] = 0$, the tangent complex $T_\pi\mathfrak{g}$ is $\mathfrak{g}[1](d_{\mathfrak{g}} + \operatorname{ad}\pi)$.

Lemma ([K], …)**.**

(i) *Let $\mathcal{U}\colon \mathfrak{g}_1^\bullet \to \mathfrak{g}_2^\bullet$ be an L_∞-map between two dg Lie algebras, $\pi \in \mathfrak{g}_1^1$ be a solution of the Maurer–Cartan equation. Then*

$$\tilde{\pi} := \mathcal{U}_1(\pi) + \frac{1}{2}\mathcal{U}_2(\pi,\pi) + \dots + \frac{1}{n!}\mathcal{U}_n(\pi,\dots,\pi) + \dots \tag{38}$$

is a solution of Maurer–Cartan equation in $\mathfrak{g}_2^\bullet$ (it is clear that $\tilde{\pi} \in \mathfrak{g}_2^1$);

(ii) *the map*

$$T_\pi\mathcal{U}\colon T_\pi\mathfrak{g}_1 \to T_{\tilde\pi}\mathfrak{g}_2,$$

defined as

$$T_\pi\mathcal{U}(x) = \mathcal{U}_1(x) + \mathcal{U}_2(x,\pi) + {} + \frac{1}{2}\mathcal{U}_3(x,\pi,\pi) + \dots + \frac{1}{(n-1)!}\mathcal{U}_n(x,\pi,\pi,\dots,\pi) + \dots \tag{39}$$

is a map of complexes.

3.5.2. In the case of the formality morphism $\mathcal{U}\colon T^\bullet_{\mathrm{poly}}(\mathbb{R}^d) \to \mathcal{D}^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ both dg algebras have an extra property: the dg spaces $T_\pi T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ and $T_{\tilde\pi}\mathcal{D}^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ have structures of associative algebras. In the case $T_\pi T^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ it is just the wedge product of polyvector fields; in the case $T_{\tilde\pi}\mathcal{D}^\bullet_{\mathrm{poly}}(\mathbb{R}^d)$ it is the usual cup-product in the deformed algebra:

$$(\Psi_1 \cup \Psi_2)(a_1 \otimes \dots \otimes a_{k+l}) = \Psi_1(a_1 \otimes \dots \otimes a_k) * \Psi_2(a_{k+1} \otimes \dots \otimes a_{k+l}). \tag{40}$$

The following remarkable result is proved in [K], Section 8:

Theorem (Kontsevich)**.** *The tangent map $T_\pi\mathcal{U}$ induces a map of the associative algebras on the level of cohomology.*

In other words, the map

$$[T_\pi\mathcal{U}]\colon H^\bullet(T_\pi T^\bullet_{\mathrm{poly}}) \to H^\bullet(T_{\tilde\pi}\mathcal{D}^\bullet_{\mathrm{poly}})$$

is a map of algebras.

Actually, Kontsevich considers a bit more general situation: a solution π of the Maurer–Cartan equation in $(\mathfrak{g}^\bullet \otimes \mathfrak{m}^\bullet)^1$ where $\mathfrak{m}^\bullet$ is a commutative dg algebra. This generality is very useful for applications, but we restrict ourselves by the case $\mathfrak{m}^\bullet = \mathbb{C}[0]$.

3.5.3. Now the tangent complex $T_{\hbar\pi}(C_\bullet(A,A)) = C_\bullet(A_*,A_*)$ has a module structure over $C^\bullet(A_*,A_*) = T_{\hbar\pi}C^\bullet(A,A)$ (for any algebra A_*):

For $\Psi\colon A_*^{\otimes k} \to A_*$ and for $\omega = a_0 \otimes a_1 \otimes \dots \otimes a_n$ $(k \le n)$ we have the "cup-product":

$$\Psi(\omega) = (a_0 * \Psi(a_1 \otimes \dots \otimes a_k)) \otimes a_{k+1} \otimes \dots \otimes a_n. \tag{41}$$

Lemma.

(i) *The "cup-product" described above*

$$\cup\colon C^\bullet(A_*,A_*) \otimes C_\bullet(A_*,A_*) \to C_\bullet(A_*,A_*)$$

is a map of the complexes;

(ii) *It endowes $C_\bullet(A_*,A_*)$ with an algebra structure over $(C^\bullet(A_*,A_*))^{\mathrm{opp}}$ (the algebra A^{opp} is the algebra with the opposite multiplication: $a *_{A^{\mathrm{opp}}} b = b *_A a$).*

Proof. It is straightforward. □

The map

$$\cup\colon T^\bullet_{\mathrm{poly}}(\mathbb{R}^d) \otimes \Omega^\bullet(\mathbb{R}^d) \to \Omega^\bullet(\mathbb{R}^d)$$

(cup-product for $\Omega^\bullet$) is defined as the operator i_γ of the insertion of the polyvector field in the differential form. It is a map of complexes:

$$\cup\colon (T^\bullet_{\mathrm{poly}}(\mathbb{R}^d), d = \operatorname{ad}\pi) \otimes (\Omega^\bullet(\mathbb{R}^d), d = L_\pi) \to (\Omega^\bullet(\mathbb{R}^d), d = L_\pi).$$

It is exactly the structure induced by (41) on the level of cohomology.

3.5.3.1.

Conjecture. *On the level of cohomology the tangent map*

$$T_{\hbar\pi}\hat{\mathcal{U}}\colon T_{\hbar\pi}C_\bullet(A,A)\to T_{\hbar\pi}\Omega^\bullet(\mathbb{R}^d)[[h]]$$

is a map of the modules, i.e. for $\omega\in T_{\hbar\pi}C_\bullet(A,A)$ *and for* $\eta\in T_{\hbar\pi}T_{\mathrm{poly}}(\mathbb{R}^d)$ *one has*:

$$[T_{\hbar\pi}\hat{\mathcal{U}}]([T_{\hbar\pi}\mathcal{U}]([\eta])\cup[\omega])=[\eta]\cup[T_{\hbar\pi}\hat{\mathcal{U}}]([\omega]) \tag{42}$$

($[\dots]$ *stands for the cohomological class of an element or for the map induced on cohomology*).

At the moment I don't know any proof of this Conjecture.

3.6. Here we consider the case of a linear Poisson structure π on $\mathbb{R}^d$, $\pi=\sum c_{ij}^k x_k\partial_i\wedge\partial_j$. Recall, that it follows from the equation $[\pi,\pi]=0$ that $\mathbb{R}^d\simeq\mathfrak{g}^*$ for a Lie algebra $\mathfrak{g}$, and the numbers $\{c_{ij}^k\}$ are the structure constants of this Lie algebra.

3.6.1. First, let us describe the map $T_\pi\hat{\mathcal{U}}|_{A=C_0(A,A)}$ in this case. We have 1 point on the circle and some points inside the disk D^2. One can prove that all possible graphs are "several wheels," as it is shown in Fig. 5.



Figure 5.

(There are no edges starting at **1** because we should obtain a 0-form, i.e. a function.) All the wheels have an even number of vertices, because of the symmetry with respect to line l (see Fig. 5). Let w_{2k} be the weight of the single wheel with $2k$ vertices (in our sense, see Section 2.2.6).

We have:

$$(T_{t\pi}\hat{\mathcal{U}})(a) = \exp\left(\sum_{k\geq 1} \hbar^{2k} w_{2k} \mathrm{Tr}_{2k}\right) a \tag{43}$$

(here $a \in S^\bullet(\mathfrak{g}) = C_0(S^\bullet(\mathfrak{g}), S^\bullet(\mathfrak{g})))$. The operator (43) satisfies the following properties:

(i) it maps $[S(\mathfrak{g})_*, S(\mathfrak{g})_*]$ to $\{S(\mathfrak{g}), S(\mathfrak{g})\}$;

(ii) if Conjecture 3.5.3.1 is true,
$$[T_{\hbar\pi}\hat{\mathcal{U}}](\omega\cdot\eta) = \omega\cdot[T_{\hbar\pi}\hat{\mathcal{U}}](\eta)$$
for $\omega \in [S(\mathfrak{g})]^{\mathfrak{g}}$ and $\omega \in S(\mathfrak{g})_*/[S(\mathfrak{g})_*, S(\mathfrak{g})_*]$.

The property (ii) follows from the "vanishing of the wheels" [Sh], according to which $T_{\hbar\pi}\mathcal{U} = \mathrm{Id}$.

Theorem. *If Conjecture* 3.5.3.1 *is true, all the numbers* w_{2k}, $k \geq 1$ *are equal to zero.*

Proof. The numbers w_{2k}, $k \geq 1$ in (43) do not depend on the Lie algebra $\mathfrak{g}$. Therefore, one can suppose that $\mathfrak{g}$ is semisimple. We proved in Section 3.1 that if $\mathfrak{g}$ is semisimple, that for $\omega \in [S^\bullet(\mathfrak{g})]^{\mathfrak{g}}$ and $\eta \in S^\bullet(\mathfrak{g})/[\mathfrak{g}, S^\bullet(\mathfrak{g})]$ one has:
$$\varphi_{D*}(\omega\cdot\eta) = \varphi_D^*(\omega)\cdot\varphi_{D*}(\eta)$$
where $\varphi_{D*}\colon [S^\bullet(\mathfrak{g})]^{\mathrm{coinv}} \to [U(\mathfrak{g})]^{\mathrm{coinv}}$ and $\varphi_D^*\colon [S^\bullet(\mathfrak{g})]^{\mathrm{inv}} \to [U(\mathfrak{g})]^{\mathrm{inv}}$ are both induced by the map $\varphi_D = \varphi_{PBW}\circ\varphi_{\mathrm{strange}}$. Next, consider the isomorphism $\Theta\colon S(\mathfrak{g})_* \to U(\mathfrak{g})$, defined as
$$\Theta(g_1 * \cdots * g_k) = g_1 \otimes \cdots * g_k.$$
It is proven in [Sh] that $\Theta = \varphi_{PBW}\circ\varphi_{\mathrm{strange}}$.

We obtain that

$$\mathrm{Id}_*(\omega\cdot\eta) = \mathrm{Id}^*(\omega)\cdot\mathrm{Id}_*(\eta) \tag{44}$$

for $\omega \in [S^\bullet(\mathfrak{g})_*]^{\mathrm{inv}}$, $\eta \in [S^\bullet(\mathfrak{g})]^{\mathrm{coinv}}$, and $\mathrm{Id}_*\colon [S^\bullet(\mathfrak{g})_*]^{\mathrm{coinv}} \to [S^\bullet(\mathfrak{g})]^{\mathrm{coinv}}$, $\mathrm{Id}^*\colon [S^\bullet(\mathfrak{g})_*]^{\mathrm{inv}} \to [S^\bullet(\mathfrak{g})]^{\mathrm{inv}}$ are induced by the identity map $\mathrm{Id}\colon S^\bullet(\mathfrak{g})_* \to S(\mathfrak{g})$. Suppose now that for a map T_* we have

$$T_*(\omega\cdot\eta) = \mathrm{Id}^*(\omega)\cdot T_*(\eta) \tag{45}$$

($T_* = T_{\hbar\pi}(\hat{\mathcal{U}})$ in our case). Then we have $T_* = \mathrm{Id}_*$. Indeed, set $\eta = [1]$ and use the decompositions
$$U(\mathfrak{g}) = Z(U(\mathfrak{g})) \oplus [U(\mathfrak{g}), U(\mathfrak{g})],$$
$$S^\bullet(\mathfrak{g}) = [S^\bullet(\mathfrak{g})]^{\mathrm{inv}} \oplus \{S(\mathfrak{g}), S(\mathfrak{g})\}$$
which hold for a semisimple Lie algebra $\mathfrak{g}$. □

3.6.2.

Corollary. *If Conjecture* 3.5.3.1 *is true, then for any Lie algebra* $\mathfrak{g}$ *Conjecture* 3.1.4 *holds for* 0*-(co)homology.*

3.6.3. Independently of Conjecture 3.5.3.1, the identity map $\mathrm{Id}\colon S^\bullet(\mathfrak{g})_* \to S^\bullet(\mathfrak{g})$ is a map of $\mathfrak{g}$-modules for any Lie algebra $\mathfrak{g}$. Indeed, the map $\varphi_D = \varphi_{PBW} \circ \varphi_{\mathrm{strange}}\colon S^\bullet(\mathfrak{g}) \to U(\mathfrak{g})$ is a map of $\mathfrak{g}$-modules, and the map $\varphi_D^{-1}\colon U(\mathfrak{g}) \to S^\bullet(\mathfrak{g})_*$ is an isomorphism of algebras (according to [Sh]). Therefore, the composition $\varphi_D^{-1} \circ \varphi_D = \mathrm{Id}$ is a map of $\mathfrak{g}$-modules. In particular, it defines a map of coinvariants $\mathrm{Id}_*\colon [S^\bullet(\mathfrak{g})_*]^{\mathrm{coinv}} \to [S^\bullet(\mathfrak{g})]^{\mathrm{coinv}}$, i.e. Id maps $[S^\bullet(\mathfrak{g})_*, S^\bullet(\mathfrak{g})_*]$ to $\{S^\bullet(\mathfrak{g}), S^\bullet(\mathfrak{g})\}$. In other words, if we set

$$[f,g]_* = \sum_{k\geq 0} \hbar^{2k+1} C_{2k+1}(f,g) \tag{46}$$

then $C_{2k+1} = \{F_{2k+1}, G_{2k+1}\} + \{F'_{2k+1}, G'_{2k+1}\} + \ldots$ for some $F_{2k+1}, G_{2k+1}, F'_{2k+1}, G'_{2k+1}, \ldots \in S^\bullet(\mathfrak{g})$. For example, $C_1(f,g) = \{f,g\}$.

It would be very interesting to calculate the map $T_{\hbar\pi}(\hat{\mathcal{U}})$ for higher cohomology and to deduce Conjecture 3.1.4 from Conjecture 3.5.3.1. The proof of Conjecture 3.5.3.1 should be somewhat very close to the proof of the Kontsevich theorem on cup-products on tangent cohomology [K], Section 8.

Finally, we have seen in this Section that the weels w_{2k}, $k \geq 1$, are conjecturally equal to zero, as well as the Kontsevich wheels [Sh]. It is very interesting to compare the Kontsevich integrals of “higher wheels“ and our integrals of them (a “higher wheel” = a graph which appear in $T_{\hbar\alpha}(\hat{\mathcal{U}})(f)$ or $T_{\hbar\alpha}(\mathcal{U})(f)$ for a non-linear α).

Acknowledgments

The first proof of the Tsygan formality conjecture for chains, based on operadic methods, appeared in the talk of Dima Tamarkin [T] at the Moshé Flato 2000 Conference. The author was inspired by this talk and by further discussions with Boris Tsygan. Many discussions with Giovanni Felder on the first version of this paper were very useful, they helped me to find the omitted in the first version boundary stratum S2.2. I thank Seva Kordonsky for a quick and quality typing of this text. I am grateful to IPDE grant 1999–2001 for the partial financial support and to the ETH-Zentrum (Zürich) where the work was done for hospitality and for the remarkable atmosphere.

References

[FSh] G. Felder, B. Shoikhet, Deformation Quantization with Traces, preprint math.QA/0002057.
[K] M. Kontsevich, Deformation quantization of Poisson manifolds, I, preprint math. q-alg/9709040.
[T] D. Tamarkin, Talk at the Conference Moshé Flato 2000, Dijon, September 11–14, 2000.
[Ts] B. Tsygan, Formality conjecture for chains, math. QA/9904132.
[Sh] B. Shoikhet, Vanishing of the Kontsevich integrals of the wheels, preprint math.QA/0007080.

FIM, ETH-Zentrum, CH-8092 Zürich, SWITZERLAND
E-mail address: `borya@mccme.ru`